

# Use of Breeding Area by Immature Spotted Eagles *Aquila pomarina* and *A. clanga* in Belarus: different strategies

V. C. Dombrovski

## ABSTRACT

During a 1999-2002 RSPB-funded APB-BirdLife Belarus project, data on the occurrence of immature individuals of both species in the breeding season were collected. 25 immature GSE and 17 immature LSE were observed. Immature GSEs were sighted with the first arriving adults (earliest date – 9 March), spending the entire breeding season in the vicinity of occupied territories in the typical GSE breeding biotopes (large forest-mire complexes), never congregating in groups. In March-early April, solitary birds in juvenile plumage were often seen accompanying an adult in its breeding area (resembling a pair from a distance). But later on, immature birds left the territories occupied by adults. Immature LSEs arrived later than immature GSEs. The earliest date was 15 April, but more than 70% of observations of immature LSEs were made in June-July, when the birds congregated in small groups of 2-5 individuals. They fed on vast drained wetlands used for hay-making. Doubtless, some several hundreds of immature LSEs stay in Belarus during summer. The low observation rate is explained only by the absence of special counts on open agricultural areas. Probably the majority of immature LSEs arrive in Belarus towards the hay-making season (late May-June). This supposition is supported by data on spring migration of two satellite-tracked immature LSEs (Meyburg *et al.* 2001).

On August 14, 2002 in Ivanovo Region (Russia), a LSE in the second plumage was sighted near the north-easternmost recorded point of the species' breeding range (Melnikov *et al.* 2001), demonstrating that imm. LSEs use the entire breeding area and are not limited to Central Europe.

## INTRODUCTION

The data on the occurrence of immature Spotted Eagles *Aquila pomarina* and *A. clanga* here presented were collected as a byproduct of counts of breeding pairs. Therefore I cannot claim to provide a comprehensive coverage of the issue. Study of the available literature, as part of the analysis of these data, has not revealed any information on the quantity and behaviour characteristics of immature Greater (GSE) and Lesser Spotted Eagles (LSE) summering in Europe. This lack of information has prompted the present article, since this facet of the population biology of Spotted Eagles is obviously understudied and it is hoped thereby to stimulate interest in the issue.

## MATERIAL AND METHODS

During the 1999-2002 RSPB-funded census of GSE and LSE, implemented by APB-BirdLife Belarus, data on the occurrence of immature individuals of both species in the breeding season were collected. Each bird's age was estimated, based on visual observation (using a 20x60 telescope) of the partly juvenile plumage and the pattern of moulting of primaries and secondaries (Forsman 1999). Throughout the period, 25 immature GSEs and 17 LSEs were sighted.

Apart from this, zoological museum collections were also examined in Moscow State University (94 GSEs and 31 LSEs) and Belarusian State University (3 GSEs and 8 LSEs).

## RESULTS AND DISCUSSION

### **Greater Spotted Eagle**

Immature GSEs arrive in Belarus with first returning adult birds. The earliest observation of immature GSEs in Olmany Marshes (Southern Belarus, around the border with Ukraine) was recorded on 9 March, 2002. In four cases, these early-arriving second calendar year (2<sup>nd</sup> cy) immature GSEs were seen together with an adult (supposedly a female, judging by its large size) around its breeding area (Table 1). At a distance, the two looked like a breeding pair. It is difficult to say whether these immature birds were offspring of these particular adults or whether they formed temporary unions. The former assumption is substantiated by the fact that, in two cases, the immature birds spent a night near the nest, with no aggression on the part of the adult bird. The above behaviour was observed only shortly after arrival (March- mid-April). Later on, immature birds were not observed in the areas occupied by the adults. Furthermore, in May-June adult birds started to show explicit aggression toward trespassing immatures.

**Table 1. Sightings of immature GSEs in Belarus during breeding season.**

<i>Date</i>	<i>Number of birds observed</i>	<i>Age of birds</i>	<i>Notes</i>
9.03.02	1	2 <sup>nd</sup> cy	Accompanying an adult female near the nest
12.03.02	2	2 <sup>nd</sup> – 3 <sup>rd</sup> cy	Migrating
27.03.02	1	2 <sup>nd</sup> cy	
1.04.00	1	2 <sup>nd</sup> cy	Near the nest of GSE
1.04.01	1	2 <sup>nd</sup> cy	Accompanying an adult female near the nest
10.04.01	1	2 <sup>nd</sup> cy	Accompanying an adult female near the nest
10.04.01	1	2 <sup>nd</sup> –4 <sup>th</sup> cy	Migrating
11.04.01	1	3 <sup>rd</sup> –4 <sup>th</sup> cy	
12.04.03	1	2 <sup>nd</sup> cy	Accompanying an adult female near the nest
20.04.99	1	2 <sup>nd</sup> cy	
20.04.00	1	2 <sup>nd</sup> cy	
21.04.00	1	2 <sup>nd</sup> – 3 <sup>rd</sup> cy	
24.04.00	1	2 <sup>nd</sup> cy	
19.05.99	1	2 <sup>nd</sup> cy	
25.06.02	1	2 <sup>nd</sup> – 3 <sup>rd</sup> cy	
25.06.02	1	3 <sup>rd</sup> –4 <sup>th</sup> cy	
27.06.02	3	2 <sup>nd</sup> – 3 <sup>rd</sup> cy	2 birds feeding together on a hayfield
5.07.01	1	3 <sup>rd</sup> –4 <sup>th</sup> cy	
15.07.00	1	2 <sup>nd</sup> cy	
15.07.02	1	3 <sup>rd</sup> –4 <sup>th</sup> cy	
15.07.01	1	3 <sup>rd</sup> –4 <sup>th</sup> cy	
16.07.99	1	2 <sup>nd</sup> cy	

Throughout the entire breeding season immature GSEs were normally observed near the areas occupied by adult birds in the species' typical habitats (vast forest wetlands, river floodplains), rarely assembling in groups. In a single case, two immature GSEs were hunting together in a hayfield like a LSE. The highest number of observations of immature GSEs was registered in the areas of high breeding density (Central Polesie in Southern Belarus). The average occurrence ratio for immature GSEs in Belarus was one bird per 5-6 breeding pairs. Thus some 25-40 immature GSEs are estimated to be present in Belarus during the breeding season.

Out of the 94 GSEs sampled in the museum of Moscow State University, 12 turned out to be immature birds of 2-4 cy, collected across the entire breeding range stretching from Europe to the Far East (Table 3).

In neighbouring Poland, isolated non-breeding GSEs (usually immature) have been sporadically recorded across the country (Tomialojc 1990). The earliest recorded date of immature GSEs in the Polish Belovezhskaya Pushcha was March 2, 1985 (Pugacewicz 1996). A winter encounter with an immature GSE in Poland was recorded on January 13, 1979 (Tomialojc 1990).

### Lesser Spotted Eagle

Immature LSEs arrive in Belarus much later than immature GSEs. Over 70% of all sightings were made in June-July (Table 2). Three observations of immature LSEs in April obviously defy the general pattern. And the earliest record of an immature LSE was on April 15, 2000 in Dikoe mire, adjacent to the National Park Belovezhskaya Pushcha, in the breeding area of a mixed pair (*Aquila pomarina x clanga*). This bird could have been the pair's previous year's offspring. In 2000, the pair raised a hybrid chick whose phenotype was that of a typical LSE (Dombrovski 2002).

In June-July immature birds were normally observed in vast agricultural areas during or directly after hay-cutting. This time of year usually sees small groups (2-5) of young and adult birds feeding peacefully side by side. At times, these hay areas are located far from forest tracts, corroborating the supposition that these groups are formed by non-breeding birds. An estimated several hundreds of immature LSEs are present in Belarus in summer. The low observation rate indicated in Table 2 can be explained by the absence of special counts on open agricultural areas during this time of year. Probably the majority of immature LSEs arrive in Belarus toward the hay-making season (late May-June). This supposition is supported by data on the spring migration of two satellite-tracked immature LSEs caught in Namibia in February 1994, one of which returned to Hungary on June 26 and the other to Ukraine on May 30 (Meyburg *et al.*, 2001).

**Table 2. Sightings of immature LSEs in Belarus during the breeding season.**

Date	Number of birds observed	Age of birds	Notes
15.04.00	1	2 <sup>nd</sup>	Near the nest of mixed pair ( <i>A.pomarina</i> x <i>A.clanga</i> )
19.04.01	1	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Migrating
20.04.01	1	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Migrating
8.06.99	1	2 <sup>nd</sup> –3 <sup>rd</sup> cy	
22.06.99	1	2 <sup>nd</sup>	
24.06.01	3	2 <sup>nd</sup> –3 <sup>rd</sup> cy	In a group of 5 LSEs feeding together in a hayfield
25.06.02	2	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Feeding together in a hayfield
1.07.00	1	2 <sup>nd</sup> –3 <sup>rd</sup> cy	In a scattered group of 10 LSEs feeding in a vast hayfield
13.07.01	1	2 <sup>nd</sup> –3 <sup>rd</sup> cy	
14.07.01	1	3 <sup>rd</sup> cy	
15.07.01	1	2 <sup>nd</sup> –3 <sup>rd</sup> cy	
19.07.00	1	2 <sup>nd</sup> –3 <sup>rd</sup> cy	
24.07.00	2	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Feeding together in a pasture

The above data challenge an account of the arrival of juvenile LSEs (younger than 4 years) in early spring (March) at breeding sites in South Bohemia (Czech Republic) (Mrlik 1998). Since adult LSEs of the South Bohemia population usually arrive very early (exceptionally the beginning of March and mid-March (Mrlik 1998)), the species in the above case could have been misidentified. Information on immature LSEs in the countries neighbouring Belarus is very scarce. A young LSE, ringed on 30 August 1956 in Lithuania, was collected on 10 July 1957 in Eastern Belarus near Orsha (R. Patapavicius, pers.comm). A 2<sup>nd</sup> cy LSE was observed by the author on August 14 2002 in the Ivanovo region of Russia (some 200km east of Moscow) near the north-easternmost recorded point of the species' breeding range (Melnikov *et al.* 2001). The bird was hunting around a cereal field.

Out of the 31 LSEs sampled in the museum of Moscow State University, 5 proved to be immature birds, collected within the breeding range, including the Moscow region of Russia (Table 3). Out of 8 LSEs in the museum of the Belarusian State University, two turned out to be immature (Table 3).

The above data indicate that in summer immature LSEs are spread across the entire breeding area of the species and not limited to Central Europe.

Interestingly, all immature LSEs in museum collections are males, though the limited size of the sample range precludes any conclusions from being drawn.

**Table 3. Immature Spotted Eagles in museum collections of Moscow and Minsk**

№	Date collected	Species	Sex	Age	Origin
<i>Moscow State University</i>					
R-37222	1.05.1904	<i>A.clanga</i>	male	2 <sup>nd</sup>	Orlov province,
R-96529	14.05.1970	-	male	2 <sup>nd</sup>	Khabarovsk Krai,
R-30361	25.05.1904	-	male	3 <sup>rd</sup> –4 <sup>th</sup> cy	Simbirsk province,
R-52720	29.05.1940	-	male	2 <sup>nd</sup>	Western Siberia,
R-30366	4.06.1904	-	female	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Simbirsk province,
R-48344	8.06.1935	-	male	2 <sup>nd</sup>	Transbaikalie,
R-55563	26.07.1911	-	?	2 <sup>nd</sup>	Western Siberia,
R-75561	26.07.1934	-	male	3 <sup>rd</sup> –4 <sup>th</sup> cy	Bashkiria, Russia
R-41491	9.08.1921	-	male	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Voronezh dstr.,
R-41483	11.08.1921	-	female	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Voronezh dstr.,
R-30364	19.08.1907	-	male	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Simbirsk province,
R-30373	22.08.1908	-	?	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Western Siberia,
R-76126	9.05.1951	<i>A.pomarina</i>	male	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Belovezhskaya
R-37201	24.05.1916	-	male	3 <sup>rd</sup> –4 <sup>th</sup> cy	Transcaucasia
R-37191	21.06.1913	-	male	2 <sup>nd</sup>	Moscow region,
R-88254	12.08.1957	-	male	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Kabardin-Balkar
R-8115	12.08.1906	-	male	2 <sup>nd</sup>	Belovezhskaya
<i>Belarusian State University</i>					
	4.06.1925	<i>A.pomarina</i>	male	2 <sup>nd</sup> –3 <sup>rd</sup> cy	Belarus
	16.07.1926	-	male	3 <sup>rd</sup> –4 <sup>th</sup> cy	Berezinsky Reserve,

Thus immature GSEs and LSEs differ in terms of time of arrival in their breeding areas, social behaviour and habitat preference. The sex distribution of immature LSEs and GSEs summering in middle latitudes might also show some discrepancy, but this assumption can only be substantiated through continued study of museum collections of the two species.

It is possible that some of the birds in partially juvenile plumage have had breeding experience, as proved for the GSE in Poland (Mejburg *et al.*, 1997).

## ACKNOWLEDGEMENTS

These data were collected as part of the APB-BirdLife Belarus project, supported by the RSPB, BirdLife International partner in the UK.

I would like to express my gratitude to Dr. Geoff Hilton (RSPB) for assisting in the organization and running of Spotted Eagle research in Belarus.

I would also like to acknowledge the contribution of my colleagues, who provided their observations of immature birds or participated in joint field surveys: D. V. Zhuravlev, L. Demongin, M. G. Dmitrenok, V. N. Melnikov, S. N. Barinov, S. V. Romanova, R. Treinys.

Particular gratitude is due to the director of Zoological Museum of Moscow State University, P. S. Tomkovich, who provided access to its sample collections, as well as to the museum staff who rendered invaluable support in sample studies. I would also like to thank the director of the Zoological Museum of the Belarusian State University, A. D. Pisanenko, for his assistance in accessing sample collections.

I also express sincere gratitude to A. E. Vintchevski for his assistance in the preparation and editing of this article.

## REFERENCES

- DOMBROWSKI, V.C 2002.** Hybridization of Lesser and Greater Spotted Eagles (*Aquila pomarina* & *A. clanga*) in Belarus: Rule or Exception? *Subbuteo* 5 (1): 23-31.
- FORSMAN, D 1999.** *The Raptors of Europe and the Middle East*. T. & A. D. Poyser London. 1999.
- MELNIKOV, V.N., D.A. SOLOVKOV, A.B. KOSTIN, N.A. EGOROVA & D.V. BOGOMOLOV 2001.** Nesting of Lesser Spotted Eagle (*Aquila pomarina*) in Ivanovo region (Russia). *Acta ornithoecol., Jena* 4.2-4 (2001): 287-289.
- MEYBURG B-U., T. MIZERA & G. MACIOROWSKI 1997.** Greater Spotted Eagle *Aquila clanga* breeding in partly juvenile plumage. *Limicola* 11: 82-87.
- MEYBURG, B-U., D.H. ELLIS, C. MEYBURG, J.M. MENDELSON & W. SCHELLER 2001.** Satellite tracking of two Lesser Spotted Eagles, *Aquila pomarina*, migrating from Namibia. *Ostrich* 72 (1&2): 35-40.
- MRLIK, V. 1998.** Spotted Eagle (*Aquila clanga*) in the Czech Republic and Lesser Spotted Eagle (*Aquila pomarina*) breeding distribution in South Bohemia, Czech Republic. *Sylvia* 34: 60-72.
- TOMIALOJC, L. 1990.** Ptaki Polski. Warszawa 1990. 462 pp.

V.C. Dombrowski  
Institute of Zoology  
27 Akademichnaya Str.  
270072 Minsk, Belarus  
E-mail: valera@biobel.bas-net.by